

Technical University of Denmark



233U/236U – A new tracer for environmental processes?

Hain, K.; Steier, P.; Eigl, R.; Froehlich, M.B.; Golser, R.; Qiao, Jixin; Quinto, F.; Sakaguchi, A.

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Hain, K., Steier, P., Eigl, R., Froehlich, M. B., Golser, R., Qiao, J., ... Sakaguchi, A. (2017). 233U/236U – A new tracer for environmental processes?. Abstract from 4th International Conference on Environmental Radioactivity, Vilnius, Lithuania.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

$^{233}\text{U}/^{236}\text{U}$ – A new tracer for environmental processes?

K. Hain¹, P. Steier¹, R. Eigl², M. B. Froehlich³, R. Golser¹, X. Hou⁴, J. Lachner¹, J. Qiao⁴, F. Quinto⁵, A. Sakaguchi²

¹Faculty of Physics, University of Vienna, Währinger Str. 17, 1090 Vienna, Austria

²Graduate School of Science, Hiroshima University, Higashi-Hiroshima, Japan

³Department of Nuclear Physics, Australian National University, Building 57, Garran Road, ACT 2601, Australia

⁴Center for Nuclear Technologies, Technical University of Denmark, DTU Risø Campus, 4000 Roskilde, Denmark

⁵Institute for Nuclear Waste Disposal, Karlsruhe Institute for Technology, Hermann-von-Helmholtz Platz 1, 76344 Eggenstein-Leopoldshafen, Germany

Keywords: $^{233}\text{U}/^{236}\text{U}$, AMS, Sellafield, thermonuclear weapon

Presenting author email: karin.hain@univie.ac.at

Apart from the production of ^{236}U by fast neutrons in thermonuclear weapons via the reaction $^{238}\text{U}(n,3n)^{236}\text{U}$, ^{236}U can be also produced in nuclear power plants and fission bombs via $^{235}\text{U}(n,\gamma)^{236}\text{U}$ using thermal neutrons. In contrast, the principle production path for ^{233}U is via the reaction $^{235}\text{U}(n,3n)^{233}\text{U}$, which requires fast neutrons with energies above 13 MeV (Gorbachev et al, 1980). Therefore, an increased production can be expected in thermonuclear weapons using Orallo (uranium enriched in ^{235}U) as blanket or tamper. Consequently, in average, fallout from nuclear weapons testings should show a higher $^{233}\text{U}/^{236}\text{U}$ ratio than emissions from thermal nuclear power plants or reprocessing plants which allows source identification for contaminations present in the environment.

However, the cross section of the reaction $^{235}\text{U}(n,3n)^{233}\text{U}$ for 14 MeV neutrons is only about 0.1 barn, as shown in Figure 1. As there is only little experimental data available for the cross-section of this reaction and the utilization of Orallo is not readily accessible for all nuclear devices which exploded during the period of atmospheric testing, the ^{233}U fallout from thermonuclear weapons can be only roughly estimated to be around one to two orders of magnitude smaller than ^{236}U fallout. Consequently, neglecting n capture on ^{232}Th in rocks and local contaminations from the ^{232}Th fuel cycle, the environmental concentrations of ^{233}U can be expected to be extremely low, so that its detection is challenging also for the highly sensitive Accelerator Mass Spectrometry (AMS).




Figure 1. Cross section in barn for the reaction $^{235}\text{U}(n,3n)^{233}\text{U}$ from model calculations (solid lines) and experimental data (blue squares) (EXFOR/ENDF, 2016)

^{236}U in environmental samples is already routinely analysed at the Vienna Environmental Research Laboratory (VERA) (e.g. Froehlich et al, 2016). VERA has recently increased its detection efficiency such that it

is now capable to detect also ^{233}U , which was demonstrated by analysing the concentration of this isotope in different types of environmental sampling material, including Irish Sea sediment, corals from the Pacific Ocean and peat bog samples from Germany. These samples were known to be affected by different contamination sources, i.e. nuclear weapons fallout and the reprocessing plant Sellafield, respectively. In average, the detected $^{233}\text{U}/^{236}\text{U}$ ratio in the environment is at a level of around 1%. However, the $^{233}\text{U}/^{236}\text{U}$ ratio in the Irish Sea sediments was around one order of magnitude lower than in the Pacific Ocean corals or the peat bog samples. These findings indicate that the $^{233}\text{U}/^{236}\text{U}$ can be indeed used for the discrimination between these two contamination sources.

In contrast to Pu, whose isotopic ratios have been successfully used for source identification in the past (Lindahl et al, 2010), uranium shows a considerably higher solubility, and thus, a conservative behaviour in water so that it can be transported over large distances (Sakaguchi et al. 2012). Being isotopes of the same element, the $^{233}\text{U}/^{236}\text{U}$ ratio is independent from the chemical behaviour in the environment as well as during sample preparation which significantly facilitates the interpretation of the measured data. For these reasons, the $^{233}\text{U}/^{236}\text{U}$ ratio could serve as a powerful tracer for environmental processes when the principle contamination sources will have been characterized.

After an introduction to possible production paths of ^{233}U and ^{236}U , respectively, first results of the $^{233}\text{U}/^{236}\text{U}$ ratio detected in samples from the different environmental reservoirs named before will be presented in this talk and the interpretation of the data will be discussed.

Gorbachev, V. M., et al. 1980. Nuclear Reactions in Heavy Elements, Pergamon Press, 1st edition
EXFOR/ENDF database, 2016

www-nds.iaea.org/exfor/servlet/X4sMakeX4

M.B. Froehlich, et al. 2016. European roe deer antlers as an environmental archive for fallout U-236 and Pu-239. J. Environ. Radioact. 151, 587-592

Sakaguchi, A., et al. 2012. Uranium-236 as a new oceanic tracer: A first depth profile in the Japan Sea and comparison with caesium-137, Earth Planet. Sci. Lett., 333-334, 165-170

Lindahl, P., et al. 2010. Plutonium isotopes as tracers for ocean processes: A review. Mar. Environ. Res., 69, 73-84.